

Amoxicillin is associated with a lower risk of further antibiotic prescriptions for lower respiratory tract infections in primary care – A database analysis spanning over 30 years

Marie Stolbrink^{a*}, Laura J Bonnett^b, John D Blakey^{c,d}

*^aInstitute of Infection and Global Health, University of Liverpool, United Kingdom;
mstolbrink@doctors.org.uk*

*^bDepartment of Biostatistics, Institute of Translational Medicine, University of
Liverpool, United Kingdom*

*^cHealth Services Research, Institute of Psychology Health and Society, University of
Liverpool, United Kingdom*

*^dRoyal Liverpool University Hospital, Royal Liverpool and Broadgreen University
Hospital NHS Trust, Prescot Road, Liverpool, United Kingdom*

Amoxicillin is associated with a lower risk of further antibiotic prescriptions for lower respiratory tract infections in primary care – A database analysis spanning over 30 years

Antibiotic prescriptions for lower respiratory tract infections occur commonly in primary care but there is uncertainty about the most effective initial treatment strategy. Both increasing antimicrobial resistance and awareness of preventable harm from medicines make resolving this uncertainty a priority. Pragmatic, real-life epidemiological investigations are needed to inform future interventional studies.

In this cross-sectional database study we analysed antibiotic prescriptions for non-pneumonic, lower respiratory tract infections (LRTI) in primary care as captured in the Optimum Care Database from 1984 to 2017. The primary outcome was a second antibiotic prescription for a LRTI code within 14 days of index prescription, the secondary outcome further antibiotic prescription for any indication. Only individuals without chronic respiratory diseases were included. We conducted univariable analysis to identify factors associated with repeat prescriptions and generate hypotheses for forthcoming projects.

We analysed 367,188 index prescriptions for LRTI. Amoxicillin was the commonest used index drug (65.1%). In 6% a second antibiotic course coded for a further LRTI was prescribed (11.2% without this coding restriction). Further antibiotic prescriptions for LRTI were significantly associated with older age, previous smoking, seven day index courses and not using amoxicillin initially. The largest effect size was seen when amoxicillin was not used as index drug (odds ratio (OR) 1.15, $p < 0.001$). This would support current prescribing practice for amoxicillin as index drug in those without respiratory disease. Prospective studies are needed to explore the observed differences.

Keywords: antibiotics; lower respiratory tract infections; primary care; amoxicillin; treatment failure; database

Introduction

Lower respiratory tract infections (LRTI) are the commonest reason for antibiotic prescriptions in primary care across Europe and the probability of antibiotic prescription for coughs and colds by general practitioners in the United Kingdom (UK) increased by 40 % between 1999 and 2011 [1,2]. It is presumed that there are variations in prescribing practice but unclear whether these result in different patient outcomes.

Primary care is an ideal target for optimising antibiotic therapy since 74 % of all UK antibiotics are prescribed here [3]. The risks of rising antimicrobial resistance and adverse effects, including infective diarrhoea, must be balanced with treatment failure [4–7].

Since real-life experiences should inform real-life decision making, we undertook this retrospective observational investigation of antibiotic prescriptions in non-pneumonic LRTI and their outcomes from 1984 to 2017. We described prescribing practice and investigated factors associated with repeat prescriptions with the aim to identify areas for further investigations.

Methods

We carried out a cross-sectional database study drawing on retrospective, electronic medical records from the Optimum Patient Care Research Database (OPCRD). We included all subjects older than 15 years who received at least one antibiotic prescription for LRTI from 27th April 1984 to 5th January 2017. Primary outcome was a new antibiotic prescription for a LRTI READ code within 14 days of initial prescription. Secondary outcome was an antibiotic prescription for any indication within 14 days of index prescription. We excluded cases with coded diagnoses of pneumonia, chronic respiratory

conditions (e.g. asthma, chronic obstructive pulmonary disease (COPD)) and those whose index antibiotic course was longer than 28 days.

The OPCRd comprised data extracted through the Optimum Patient Care (OPC) Clinical Service Evaluation (<http://optimumpatientcare.org/opcrd/>). The OPCRd is a research-quality primary care database with a focus on respiratory diseases. It contained anonymised, routinely recorded patient data from over 525 UK general practices. The OPCRd was approved by the Trent Multi-Centre Research Ethics Committee for clinical research use. The study protocol was approved by the OPCRd's Anonymised Data Ethics Protocols and Transparency Committee.

Data was analysed using SPSS version 22 (IBM). Univariable analysis was carried out for gender, age, body mass index (BMI), smoking status, index antibiotic drug and index duration. Statistically significant results were defined as $p < 0.05$, 95% confidence intervals were given where appropriate. Summary statistics, odds ratios, Chi-squared and student T-tests were presented as appropriate for each variable based on type and distribution of data. Subgroup analysis was carried out for cases with more than one index prescription and cases from 1st January 2011 to 5th January 2017 to establish if the observed effects were maintained in current antibiotic prescribing practice.

Results

There were 1,549,402 antibiotic prescriptions in the OPCRd from 1984-2017; of these 753,885 were for "simple" LRTI (ie no underlying asthma, COPD or other chronic lung disease) in patients over 15 years. "Chest infection not otherwise specified", "bronchitis" and "lower respiratory tract infections" were the commonest codes for LRTI (67.5%, 21.7%, 10.2% respectively). We analysed the 367,188 cases (51%) for which an index duration was clearly documented (see supplementary figure).

1
2 Of the total eligible 367,188 index prescriptions, 58.9% occurred in female
3
4 patients. Mean age was 55.4 years, range 16-106 years. Mean BMI was 28.1 ± 6.1
5
6 (standard deviation, SD). 35.4% were non-smokers, 27% current smokers and 28.9%
7
8 were ex-smokers (missing smoking status in 8.7%). Most prescriptions occurred from
9
10 2006-2010 (35.3%). Most patients received only one index prescription (59.3%), with
11
12 20.4% having two and 20.3% three or more separate index prescriptions. The commonest
13
14 index duration was 7 days (74.1%), followed by 5 or 6 days (22.7%). Amoxicillin was
15
16 the index antibiotic prescribed most frequently (65.1%), clarithromycin and erythromycin
17
18 were the next most frequent (9.1% and 7.4% respectively). 548 cases received 2 further
19
20 antibiotic courses within 14 days of the index prescription – these were excluded from
21
22 further analysis.
23
24
25
26
27
28
29
30
31
32

33 ***Further antibiotic prescriptions for LRTI***

34

35
36 41,227 cases had a second antibiotic prescription within 14 days of the index prescription
37
38 (11.2%). More than half of the additional antibiotic prescriptions were coded for another
39
40 LRTI (54.8%, 22,176 of 41,227). Most of the 22,176 second prescriptions for another
41
42 LRTI code were for 7 days (75.2%). Clarithromycin, amoxicillin and doxycycline were
43
44 the most commonly used second line antibiotics (23.3%, 15.5% and 15.1% respectively).
45
46
47

48 Those receiving further antibiotics for LRTI were significantly older (mean 57.5
49
50 versus 54.1 years, mean difference -3.4 (MD), 95% CI -3.6 - -3.1, $p < 0.001$) than those
51
52 not receiving antibiotics (see table 1). Current smokers had a significantly reduced risk
53
54 of repeat antibiotic prescription for LRTI compared to non-smokers (OR 0.78, 95% CI
55
56 0.78-0.81, $p < 0.001$). Ex-smokers were more likely to receive repeat antibiotics than
57
58 non-smokers (OR 1.04, 95% CI 1.01-1.07, $p = 0.02$). Ex-smokers were at significantly
59
60
61
62
63
64
65

1 higher risk of repeat antibiotic prescription compared to current smokers (OR 1.34, 95%
2 CI 1.29-1.39, $p<0.001$). Not using amoxicillin as index antibiotic was associated with
3 more repeat antibiotic prescriptions (OR 1.15, 95% CI 1.11-1.18, $p<0.001$). Seven day
4 index courses were associated with more repeat prescriptions than both shorter and
5 longer courses (OR 0.93, 95% CI 0.90-0.96, $p<0.001$; OR 0.86, 95% CI 0.79-0.94,
6 $p=0.001$ respectively). Higher BMI was associated with further prescriptions (mean
7 28.5 ± 6.3 versus 28.1 ± 6.7 , MD -0.46, 95% CI -0.6 - -0.4, $p<0.001$). There was no
8 difference gender ($p=0.08$).

9
10
11
12
13
14
15
16
17
18
19
20 **[Table 1 here]**

21
22
23 Figure 1 shows that most repeat antibiotic prescriptions occurred following an
24 index duration of 7 days. The proportion of cases receiving further antibiotics decreased
25 with longer and shorter index duration. Fewer repeat prescriptions occurred when
26 amoxicillin was used as index antibiotic.

27
28
29
30
31
32 **[Figure 1 here]**

33 34 35 36 *Subgroup analysis for repeat LRTI code from 2011-2017*

37 89,694 cases were included in this period. The most commonly prescribed index duration
38 was again 7 days (83.7%). The most frequently used index antibiotics were amoxicillin
39 (70.3%), clarithromycin (14.7%) and co-amoxiclav (4.8%).

40
41
42
43
44
45
46 Men were less likely to receive further antibiotics (OR 0.93, $p=0.01$, see
47 supplementary table 2). Those receiving antibiotics were significantly older (mean age
48 60 versus 57 years, $p<0.001$) with a higher BMI (29.0 versus 28.6, $p<0.001$). Current
49 smokers were less likely to receive antibiotics (OR 0.81, 95% CI 0.76-0.87, $p<0.001$)
50 than non-smokers, ex-smokers more likely than non-smokers (OR 1.05,
51 95% CI 0.99-1.11, $p=0.11$) and significantly more likely than current smokers (OR 1.29,
52
53
54
55
56
57
58
59
60
61
62
63
64
65

95% CI 1.20-1.38, $p<0.001$). Not using amoxicillin as index antibiotic retained a higher risk of repeat prescription (OR 1.11, 95% CI 1.04-1.17, $p<0.001$). In contrast to the overall study findings shorter durations were now associated with a higher risk of repeat prescription than 7 day courses (OR 1.12, 95% CI 1.05-1.20, $p=0.001$). Longer durations maintained their lower risk of repeat antibiotics (OR 0.85, 95% CI 0.71-1.01, $p=0.06$).

Further antibiotic prescription for any indication

When considering all antibiotic prescriptions within 14 days of index prescription (41,227 cases) clarithromycin, amoxicillin and doxycycline were used most commonly as second line agents (20.5%, 17.0% and 13.7%).

Those receiving further courses were significantly older (mean 57.2 versus 54.0 years, $p<0.001$) with a significantly higher BMI (mean 28.5 versus 28.1, $p<0.001$, see supplementary table 3). Men were significantly less likely to receive antibiotics than women (OR 0.93, 95% CI 0.91-0.95, $p<0.001$). Current smokers were less likely, ex-smokers more likely to receive antibiotics than non-smokers (OR 0.79, 95% CI 0.77-0.82, $p<0.001$ and OR 1.04, 95% CI 1.01-1.07, $p=0.002$ respectively). Ex-smokers were significantly more likely to receive further antibiotics than current smokers (OR 1.31, 95% CI 1.28-1.35, $p<0.001$). Not using amoxicillin as index drug was associated with higher risk of repeat antibiotics (OR 1.18, 95% CI 1.16-1.21, $p<0.001$; see figure 1b). Shorter index courses were associated with fewer repeat prescriptions than 7 day index courses (OR 0.94, 95% CI 0.91-0.96, $p<0.001$). There was no significant difference between longer and 7 day courses (OR 0.94, 95% CI 0.88-1.00, $p=0.06$).

Discussion

We present a report of antibiotic prescribing for LRTI in over 525 primary care practices in the United Kingdom over 33 years. More than 365,000 truly representative, index prescriptions were analysed and more than one in twenty received two antibiotic courses for LRTI within 14 days. Our hypothesis-generating findings support the widespread use of amoxicillin as index drug and suggest that initial courses shorter than seven days could potentially be effective, as they were associated with a lower risk of repeat prescriptions. These findings should be confirmed in interventional studies.

A key output from this study is a description of the frequency of and variation in prescribing practice for LRTI in primary care. Repeat antibiotic prescribing was common and in discrepancy to UK guidelines: the National Institute for Health and Care Excellence does not recommend routine antibiotic prescription for those presenting with likely self-limiting illnesses, including acute bronchitis, in primary care since in the majority of cases no pathogen or only respiratory viruses can be identified [8,9]. Yet 6.2% of our population even received two antibiotic courses for LRTI within two weeks. Treatment failure rates of 20% have been described previously but these included prescriptions for pneumonia and further antibiotics for all indications [10]. We present more specific data for LRTI, which can assist to plan future work. Most cases had only one index prescription, however, subgroup analysis of those with more index prescriptions showed no significant differences to the presented findings.

Our findings were inconclusive regarding duration, highlighting the need for interventional research. Overall, shorter and longer courses than seven days seemed to be more successful at reducing the need for further antibiotics for LRTI. Shorter courses also had a lower risk for all repeat indications. However, shorter courses were associated with increased repeat prescriptions in the subgroup analysis from 2011-2017, whilst longer courses maintained their superiority over seven day courses. Yet for all indications there

1 was no difference between longer and seven day courses. With our data we are unable to
2 establish whether this reflects changes in pathogens and disease or changes in presenting
3 and prescribing behaviour by patients and clinicians. We did not have access to adverse
4 effect data, which would be important for any future studies. A recent meta-analysis
5 found an association of adverse effects and longer antibiotic duration in COPD patients
6 [11].
7
8
9
10
11
12
13

14 Repeat prescriptions were more common in older subjects, those with a higher
15 BMI or previous smoking history. These findings appear to be clinically relevant and
16 plausible as e.g. older patients are at higher risk of adverse outcomes from community
17 acquired pneumonia [9]. Patients with previous smoking history are at risk of abnormal
18 lung architecture, COPD and hence atypical bacterial infections [12]. Higher BMI may
19 impair clearing of secretions but the mean difference in BMI was only 0.4 between groups
20 which is unlikely to relate to clinical significance. Further work should be undertaken as
21 these groups may have different needs for drugs and durations for LRTI due to co-
22 morbidities and different lung architecture. Amoxicillin treats *Streptococcus*
23 *pneumoniae*, the most common bacterial cause of LRTI, hence our finding of fewer repeat
24 prescriptions after initial amoxicillin therapy could be clinically plausible [9]. Whilst the
25 effect sizes were small, LRTI are very common [9]. Hence even small differences of less
26 than 1% could theoretically result in meaningful differences in antibiotic prescriptions
27 due to the size of the population, if they were true. It may be worth exploring this in the
28 future.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Current smokers seemed to have a significantly reduced risk of repeat
52 prescriptions (OR 0.78, 95% CI 0.75-0.81). Doctors may attribute cough in a current
53 smoker to the smoking and hence not prescribe antibiotics. This is in contrast to
54 retrospective European data where current smoking was an independent risk factor for
55
56
57
58
59
60
61
62
63
64
65

1 antibiotic prescription in primary care patients presenting with cough and the increased
2 risk of community acquired pneumonia and death of pneumococcal disease in smokers
3 [11–13]. Yet the cough study demonstrated no improved recovery with antibiotics and
4
5 subgroup analysis of a multi-centre randomised, placebo-controlled trial of amoxicillin
6
7 versus placebo in LRTI showed no clinically meaningful advantage of antibiotic
8
9 treatment, supporting our described UK findings [14].
10
11
12
13

14 We analysed electronic health records. There is a drive to use more routinely
15 collected data, particularly in primary care research, e.g. from the National Institute of
16 Health Research, and highlighted by recent successes such as the Salford Lung Study, to
17 reduce burden on patients, bias and approximate real-world practice [13,14]. However,
18 the reliance on primary care data is also this study's major limitation – specifically the
19 49% of cases for which no index antibiotic duration was documented. The excluded cases
20 had a lot of missing data (see supplementary table 4). The missing data had fewer non-
21 smokers (8.8%) but also more missing smoking data (26%). It is unclear whether this
22 represents a true difference in prevalence of non-smoking or failure to code an absence
23 of disease. We were unable to comment on the repeat prescriptions in the non-included
24 group due to too much missing data. One may suspect that documentation improved over
25 time with advancing information technology, however, from 2011 to 2017 there were
26 206,455 cases in the database – and only 89,694 with a documented index duration (43%).
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Other limitations include lack of information on other healthcare contacts or signs
of infection and on other confounders, e.g. non-respiratory co-morbidities, allergies, prior
antibiotic use, unattainability of total number of LRTI in the study period and inability to
distinguish appropriate from inappropriate prescribing. Yet treatment decisions are often
made independent of these, e.g. initial clinical severity scores and clinical response to

1 therapy were not related to duration of antibiotic therapy in adults admitted with
2 community acquired pneumonia [15]. Variations in clinical presentation did not explain
3 the large discrepancies in antibiotic prescribing for acute cough in a cross-sectional study
4 of 14 European countries [16]. Repeat prescriptions could have been issued due to
5 genuine anti-microbial failure, but also due to drug failure caused by side effects or
6 incorrect diagnoses. Better data is needed to distinguish these.
7
8
9
10
11
12
13

14 Nevertheless, we present a hypothesis-generating real-world picture of the
15 prescribing patterns of antibiotics for LRTI over 33 years in 7% of all UK primary care
16 practices, analysing over 365,000 cases. With evidence on effective antibiotic treatment
17 currently lacking it is paramount to have a historical, pragmatic perspective as a basis to
18 design safe interventional studies in the future.
19
20
21
22
23
24
25
26
27
28
29
30

31 **Conclusion**

32 Antibiotics were prescribed for LRTI in people without chronic chest conditions. There
33 was variation in practice and apparent deviation from guidelines. Amoxicillin as index
34 drug seemed more successful in avoiding repeat antibiotic prescriptions than other agents,
35 supporting the demonstrated current prescribing practice. The results for optimal
36 antibiotic duration were less clear, but suggested that shorter index durations may be
37 appropriate for LRTI. This pragmatic study has implications for further work
38 around guideline implementation, antimicrobial resistance strategies,
39 and interventional studies of antibiotic duration.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 **Acknowledgements**

57 Derek Skinner, data analyst, and Victoria Carter, Optimum Patient Care, Cambridge,
58
59
60
61
62
63
64
65

UK; Alison Chisholm, previously at Optimum Patient Care, Cambridge, UK. In collaboration with the Respiratory Effectiveness Group.

Declaration of interest statement

We are grateful to the Eleanor Peel Trust for providing funding for bespoke data extraction and database creation from the OPC. Marie Stolbrink is funded by a National Institute for Health Research Academic Clinical Fellowship. Laura J Bonnett is funded by a NIHR Post-Doctoral Fellowship (PDF-2015-08-044). The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research, or the Department of Health.

References

1. Hawker JI, Smith S, Smith GE, et al. Trends in antibiotic prescribing in primary care for clinical syndromes subject to national recommendations to reduce antibiotic resistance, UK 1995-2011: analysis of a large database of primary care consultations. *J Antimicrob Chemother* [Internet]. 2014 [cited 2016];69:3423–3430. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25091508>
2. Goossens H, Ferech M, Vander Stichele R, et al. Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *Lancet*. 2005;365:579–587.
3. Public Health England. English surveillance programme for antimicrobial utilisation and resistance (ESPAUR) - Executive summary and recommendations extracted from 2016 report [Internet]. 2016. Available from: <https://www.gov.uk/government/publications/english-surveillance-programme-antimicrobial-utilisation-and-resistance-espaur-report>
4. World Health Organization. Global action plan on antimicrobial resistance

[Internet]. 2015 [cited 2015]. Available from: <http://www.who.int/antimicrobial-resistance/publications/global-action-plan/en/>

5. Ashiru-Oredope D, Sharland M, Charani E, et al. Improving the quality of antibiotic prescribing in the NHS by developing a new Antimicrobial Stewardship Programme: Start Smart--Then Focus. *J Antimicrob Chemother* [Internet]. 2012 [cited 2015];67 Suppl 1:i51-63. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22855879>
6. Dubberke ER, Carling P, Carrico R, et al. Strategies to Prevent *Clostridium difficile* Infections in Acute Care Hospitals: 2014 Update. *Infect Control Hosp Epidemiol* [Internet]. Cambridge University Press; 2014 [cited 2017];35:S48–S65. Available from: https://www.cambridge.org/core/product/identifier/S0899823X00193857/type/journal_article
7. Smith Susan M, Fahey T, Smucny J, et al. Antibiotics for acute bronchitis [Internet]. *Cochrane Database Syst. Rev.* John Wiley & Sons, Ltd; 2014. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD000245.pub3/abstract>
8. National Institute for Health and Care Excellence. Respiratory tract infections (self-limiting): prescribing antibiotics. *Clin Guidel. NICE*; 2008;
9. Currie CJ, Berni E, Jenkins-Jones S, et al. Antibiotic treatment failure in four common infections in UK primary care 1991-2012: longitudinal analysis. *BMJ* [Internet]. British Medical Journal Publishing Group; 2014 [cited 2018];349:g5493. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25249162>
10. Stolbrink M, Amiry J, Blakey JD. Does antibiotic treatment duration affect the outcomes of exacerbations of asthma and COPD? A systematic review. <https://doi.org/10.1177/1479972317745734> [Internet]. SAGE PublicationsSage UK: London, England; 2017 [cited 2017]; Available from: <http://journals.sagepub.com/doi/full/10.1177/1479972317745734#articleCitationDownloadContainer>
11. Stanton N, Hood K, Kelly MJ, et al. Are smokers with acute cough in primary care prescribed antibiotics more often, and to what benefit? An observational

study in 13 European countries. *Eur Respir J*. 2010;35.

12. Torres A, Peetermans WE, Viegi G, et al. Risk factors for community-acquired pneumonia in adults in Europe: a literature review. *Thorax*. 2013;68:1057–1065.
13. Bello S, Menéndez R, Antoni T, et al. Tobacco smoking increases the risk for death from pneumococcal pneumonia. *Chest*. 2014;146:1029–1037.
14. Moore M, Stuart B, Coenen S, et al. Amoxicillin for acute lower respiratory tract infection in primary care: subgroup analysis of potential high-risk groups. *Br J Gen Pract* [Internet]. 2014 [cited 2015];64:e75-80. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3905438&tool=pmcentrez&rendertype=abstract>
15. “NIHR Statistics Group.” NIHR Statistics Group Annual Meeting [Internet]. *Summ. Annu. Gen. Meet.* 2017 [cited 2018]. Available from: <https://statistics-group.nihr.ac.uk/nihr-statistics-group-annual-meeting-challenges-and-opportunities-for-applied-statisticians/>
16. New JP, Bakerly ND, Leather D, et al. Obtaining real-world evidence: the Salford Lung Study. *Thorax* [Internet]. 2014;69:1152–1154. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4251297/>
17. Aliberti S, Blasi F, Zanaboni AM, et al. Duration of antibiotic therapy in hospitalised patients with community-acquired pneumonia. *Eur Respir J* [Internet]. 2010 [cited 2016];36:128–134. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19926738>
18. Butler CC, Hood K, Verheij T, et al. Variation in antibiotic prescribing and its impact on recovery in patients with acute cough in primary care: prospective study in 13 countries. *BMJ* [Internet]. *BMJ Group*; 2009 [cited 2017];338:b2242. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19549995>

Tables with captions

Table 1: Univariable analysis for repeat antibiotic prescription with LRTI code (total number 366,640)

	Receiving second antibiotic course for LRTI			
Analysed variable	Yes	No	Calculation	p value
Age (years)	57.5 ± 16.6	54.1 ± 17.7	MD -3.4 (95% CI -3.6 - -3.1)	< 0.001
Smoking status				
Non-smoker	8,528 (6.6%)	121,363 (93.4%)	OR 1.00	
Current smoker	5,135 (5.2%)	93,929 (94.8%)	OR 0.78 (95% CI 0.75-0.81)	< 0.001
Ex-smoker	7,208 (6.8%)	98,737 (93.2%)	OR 1.04 (95% CI 1.01 – 1.07)	0.02
Missing			n = 31,740	
Index drug				
Amoxicillin	13,905 (5.8%)	224,716 (94.2%)	OR 1.00	
Not amoxicillin	7,323 (6.6%)	103,339 (93.4%)	OR 1.15 (95% CI 1.11-1.18)	< 0.001
Missing			n = 17,357	
Index antibiotic duration				
7 days	16,732 (6.2%)	254,769 (93.8%)	OR 1.00	
< 7 days	4,904 (5.8%)	80,188 (94.2%)	OR 0.93 (95% CI 0.90-0.96)	< 0.001
> 7 days	540 (5.4%)	9,507 (94.6%)	OR 0.86 (95% CI 0.79-0.94)	0.001
BMI (kg/m²)	28.5 ± 6.3	28.1 ± 6.7	MD -0.46 (95% CI -0.6 - -0.4)	< 0.001
Gender				
Female	12957 (6.1%)	199,185 (93.9%)	1.00	0.08
Male	9219 (6.0%)	145,279 (94.0%)	OR 0.98 (95% CI 0.95-1.00)	

(Mean ± SD or absolute number + row % where appropriate; SD: standard deviation;

OR: odds ratio; 95% CI: 95% Confidence interval; MD: mean difference; p value as per students T-test or Chi-Square)

Figure 1





Figure captions:

Figure 1: Percentage of all prescriptions of specified index duration and index antibiotic receiving second antibiotic prescription for LRTI (a) or any indication (b) within 14 days of index antibiotic prescription, by index duration and antibiotic group. (* $p < 0.005$ by Chi-square test)

Figure 1a






Figure 1b



Supplementary data



Supplementary Figure 1

Flowchart of cases included from the OPCRD

Supplementary Table 1

Supplementary Table 1: Baseline characteristics (total number 367,188)

Characteristic	N (%) or mean \pm SD
<u>Gender</u>	
Female	216,273 (58.9%)
Male	150,915 (41.1%)
<u>Age</u>	55.4 \pm 18.0
<u>BMI</u>	28.1 \pm 6.1
<u>Smoking status</u>	
Non-smoker	130,109 (35.4%)
Current smoker	99,188 (27.0%)
Ex-smoker	106,210 (28.9%)
Missing	31,771 (8.7%)
<u>Number of index prescriptions per patient</u>	
1	217,742 (59.3%)
2	74,907 (20.4%)
≥ 3	74,539 (20.3%)
<u>Year of index prescription[#]</u>	
Earlier and up to 2000	60,324 (16.4%)
2001-2005	87,532 (23.8%)
2006-2010	129,463 (35.3%)
Since 2011	89,869 (24.5%)
<u>Index duration</u>	
≤ 4 days	1,704 (0.5%)
5 and 6 days	83,477 (22.7%)
7 days	271,955 (74.1%)
8-10 days	7,188 (2.0%)
≥ 11 days	2,864 (0.7%)
<u>Index antibiotic</u>	
Amoxicillin	238,948 (65.1%)
Clarithromycin	33,307 (9.1%)
Erythromycin	27,097 (7.4%)
Co-amoxiclav	13,862 (3.8%)
Cefalexin	12,573 (3.4%)
Doxycycline	6,302 (1.7%)
Cefaclor	6,274 (1.7%)
Ciprofloxacin	6,087 (1.7%)
Oxytetracycline	4,136 (1.1%)
Ampicillin	1,218 (0.3%)
Missing	17,384 (4.7%)

(#one individual can be included in multiple periods)

Supplementary Table 2

Univariable analysis for repeat LRTI code from 2011-2017 (total number = 89,694)

	Receiving second antibiotic course for LRTI			
Analysed variable	Yes	No	Calculation	p value
<u>Age</u> (years)	60.3 ± 17.3	57.3 ± 18.5	Mean difference: -3.0 (95 % CI -3.4 - -2.5)	<0.001
<u>Smoking status</u>				
Non-smoker	3,077 (7.6 %)	37,297 (92.4 %)	OR 1.00	
Current smoker	1,301 (6.3 %)	19,359 (93.7 %)	0.81 (95 % CI 0.76-0.87)	< 0.001
Ex-smoker	2,112 (8.0 %)	24,410 (92.0 %)	1.05 (95 % CI 0.99-1.11)	0.11
Missing			2,138 (2.4 %)	
<u>Index drug</u>				<0.001
Amoxicillin	4,497 (7.1 %)	58,538 (92.9 %)	OR 1.00	
Not amoxicillin	2,009 (7.8 %)	23,661 (92.2 %)	OR 1.11 (95 % CI 1.04-1.17)	
Missing			989 (1.1 %)	
<u>Index antibiotic duration</u>				
7 days	5,433 (7.2 %)	69,672 (92.8 %)	1.00	
< 7 days	994 (8.1 %)	11,352 (91.9 %)	1.12 (95 % CI 1.05-1.20)	0.001
> 7 days	139 (6.2 %)	2,104 (93.8 %)	0.85 (95 % CI 0.71-1.01)	0.06
<u>BMI (kg/m²)</u>	29.0 ± 6.5	28.6 ± 6.4	Mean difference: -0.4 (95 % CI -0.6 - -0.3)	<0.001
<u>Gender</u>				0.009
Female	3,847 (7.5 %)	47,330 (92.5 %)	OR 1.00	
Male	2,719 (7.1 %)	35,798 (92.9 %)	OR 0.93 (95 % CI 0.89 – 0.98)	

Supplementary Table 3

Univariable analysis for repeat antibiotic prescription for any indication within 14 days (total number = 366,640)

	Receiving second antibiotic course for any indication			
Analysed variable	Yes	No	Calculation	p value
<u>Age</u> (years)	57.2 ± 16.8	54.0 ± 17.7	Mean difference: -3.2 (95 % CI -3.4 - -3.1)	<0.001
<u>Smoking status</u>				
Non-smoker	15,418 (11.9 %)	114,473 (88.1 %)	OR 1.00	
Current smoker	9,564 (9.7 %)	89,500 (90.3 %)	OR 0.79 (95 % CI 0.77-0.82)	< 0.001
Ex-smoker	13,022 (12.3 %)	92,923 (87.7 %)	OR 1.04 (95 % CI 1.01-1.07)	0.002
Missing			31,740 (8.7 %)	
<u>Index drug</u>				<0.001
Amoxicillin	25,536 (10.7 %)	213,085 (89.3 %)	OR 1.00	
Not amoxicillin	13,752 (12.4 %)	96,910 (87.6 %)	OR 1.18 (95 % CI 1.16-1.21)	
Missing			17,357 (4.7 %)	
<u>Index antibiotic duration</u>				
7 days	30,977 (11.4 %)	240,524 (88.6 %)	OR 1.00	
< 7 days	9,164 (10.8 %)	75,928 (89.2 %)	OR 0.94, (95 % CI 0.91-0.96,	< 0.001
> 7 days	1,086 (10.8 %)	8,961 (89.2 %)	OR 0.94, (95 % CI 0.88-1.00,	0.06
<u>BMI (kg/m²)</u>	28.5 ± 6.3	28.1 ± 6.8	Mean difference: -0.4 (95 % CI -0.5 - -0.3)	<0.001
<u>Gender</u>				<0.001
Female	24,510 (11.6 %)	187,632 (88.4 %)	OR 1.00	
Male	16,717 (10.8 %)	137,781 (89.2 %)	OR 0.93 (95 % CI 0.91-0.95)	

Supplementary table 4

Comparison of characteristics of included and excluded patients based on documentation of index antibiotic duration

Characteristic	Included patients (n=367,188) N (%) or mean \pm SD	Excluded patients (n=386,697) N (%) or mean \pm SD
<u>Gender</u>		
Female	216,273 (58.9%)	213,244 (59.8%)
Male	150,915 (41.1%)	173,453 (40.2%)
<u>Age</u>	55.4 \pm 18.0	56.3 \pm 18.1
<u>BMI</u>	28.1 \pm 6.1	28.1 \pm 6.3
<u>Smoking status</u>		
Non-smoker	130,109 (35.4%)	34,029 (8.8%)
Current smoker	99,188 (27.0%)	119,489 (30.9%)
Ex-smoker	106,210 (28.9%)	132,637 (34.3%)
Missing	31,771 (8.7%)	100,542 (26.0%)
<u>Number of index prescriptions per patient</u>		
1	217,742 (59.3%)	110,209 (28.5%)
2	74,907 (20.4%)	80,433 (20.8%)
≥ 3	74,539 (20.3%)	196,055 (50.7%)
<u>Year of index prescription[#]</u>		
Earlier and up to 2000	60,324 (16.4%)	Not documented
2001-2005	87,532 (23.8%)	Not documented
2006-2010	129,463 (35.3%)	Not documented
Since 2011	89,869 (24.5%)	Not documented
<u>Index duration</u>		
≤ 4 days	1,704 (0.5%)	Not documented
5 and 6 days	83,477 (22.7%)	Not documented
7 days	271,955 (74.1%)	Not documented
8-10 days	7,188 (2.0%)	Not documented
≥ 11 days	2,864 (0.7%)	Not documented
<u>Index antibiotic</u>		
Amoxicillin	238,948 (65.1%)	235,498 (60.9%)
Not amoxicillin	110,308 (30.2%)	132,250 (34.2%)
Missing	17,384 (4.7%)	18,949 (4.9%)
<u>Repeat antibiotics for LRTI</u>	(n=366,640)	
Yes	22,176 (6.0%)	Not documented
No	345,012 (94.0%)	Not documented
<u>Repeat antibiotics for all indications</u>	(n=366,640)	
Yes	41,227 (11.2%)	Not documented
No	325,961 (88.8%)	Not documented

<u>Repeat antibiotics for LRTI code 2011-2017</u>	(n=89,694)	
Yes	6,566 (7.3%)	Not documented
No	83,128 (92.7%)	Not documented